

Care recommendations for the respiratory complications of esophageal atresia-tracheoesophageal fistula

This summary document is based on a publication by Koumbourlis et al.¹ and incorporates recent statements and guidelines of the European Respiratory Society as well as consensus recommendations of INoEA, ERNICA and ERN-LUNG working groups.

Esophageal atresia (EA) and tracheoesophageal fistula (TEF) present significant lifelong respiratory challenges. Involvement of a physician with expertise in respiratory care is therefore recommended, specifically in infancy and at the preschool age. The recommendations of the INoEA's Respiratory Complications Working Group (RCWG)¹ provide a structured approach to the diagnosis, treatment, and long-term care of these complications, aiming to improve patient outcomes. The RCWG identified the primary respiratory conditions faced by patients before and/or after EA-TEF repair, including:

1. Tracheobronchomalacia (TBM)





Figure 1) Normal trachea during bronchoscopy on the left, severe tracheomalacia (>75% of lumen narrowing) shown on the right. The degree of obstruction may vary among patients.

TBM is the partial or complete collapse of the main windpipe (trachea and/or main stem bronchi) due to abnormal airway wall softness or floppy cartilage. It is present in nearly all EA-TEF patients, with severity varying based on airway narrowing.



TBM may cause acute symptoms in neonates and infants but can also result in long-term complications.

Symptoms of TBM according to age		
Neonates-Infants	Children-Adolescents	Adults
Respiratory distress	Recurrent wheeze	Recurrent wheeze
Apnea	"Honking"/"Brassy" and/or "wet" cough	Recurrent "wet" and/or "honky" cough
Episodes of severe hypoxemia (low oxygen saturation or "blue" or "dusky" spells)	Hoarse voice/Stridor	Hoarse voice
Recurrent harsh "wheeze"	Exercise Intolerance	Exercise intolerance
"Honking"/"Brassy" cough	Obstructive lung function, low peak expiratory flow (PEF)	Obstructive lung function, low peak expiratory flow (PEF)
Stridor/"hoarse" cry		
Recurrent airway infections	Recurrent airway infections	Recurrent airway infections

The symptoms of TBM vary according to the age of the patient. They tend to be most severe in the neonatal period and generally may improve with time.

Diagnosis of TBM: Flexible bronchoscopy under light general anesthesia with spontaneous breathing without positive end expiratory pressure (PEEP) is the most informative method for assessing TBM presence, extent, and severity. Dynamic CT scans and advanced MRIs can also aid evaluation. The shape of the flow volume curve from spirometry can be indicative of TBM.




Figure 2) Example of flow-volume curves during forced expiration before (pre) and after (post) inhalation of a short acting beta-mimetic in a patient with esophageal atresia and tracheo-bronchial malacia. In this patient, there was no major change with bronchodilator. Normal range is shown in the shaded grey area. In some cases, beta-mimetics have a negative effect on expiratory flow. Curves vary between patients.

Management of TBM: Interventions aim to reduce symptoms due to TBM and, when it is severe, reduce TBM severity by surgery. In neonates, the primary goal is to prevent respiratory distress and low oxygen levels. Treatments range from supplemental oxygen use to non-invasive positive airway pressure. Severe cases may require intubation, mechanical ventilation, or surgery (tracheopexy/aortopexy) generally after the initial EA repair. Rarely, a tracheostomy is needed.

Long-term management of TBM focuses on clearing airway secretions and preventing airway infections. This can be achieved through age-appropriate airway clearance techniques (in adults: active cycle of breathing techniques), positive expiratory pressure devices such as PEP masks or oscillatory positive pressure devices^{2,3}. Nebulized hypertonic saline may help to improve mucociliary clearance and inhaled corticosteroids may reduce airway inflammation in selected patients. Infections, often signaled by a "wet" cough or fever, are treated aggressively with antibiotics. Bronchodilators should be used cautiously, as they can worsen airway collapse.

2. Aspiration Risk

Aspiration Risk in EA-TEF: EA-TEF patients are highly prone to aspiration due to multiple factors:

- Esophageal dysmotility and/or strictures, causing food or fluids to enter the airway.
- Recurrent TEF.
- High prevalence of gastroesophageal reflux (GER).
- Occasionally, a laryngeal cleft.

Diagnosis: A systematic approach is essential to evaluate all potential aspiration causes. Swallowing function can be assessed via videofluoroscopic or modified contrast swallow studies. Esophageal narrowing and function are diagnosed by contrast swallow studies and sometimes with manometry. Recurrent TEF is diagnosed using a "pull-back" contrast study, bronchoscopy and/or upper endoscopy. GER is identified clinically or through contrast studies, pH-impedance monitoring, or endoscopy. Chronic bacterial infection due to aspiration can be detected through repeated sputum cultures and chest imaging, and if symptoms persist despite optimal treatment by bronchoscopy with bronchoalveolar lavage (BAL) to determine pathogens. Collaboration with a gastroenterologist otolaryngologist, occupational therapist, speech therapist, physiotherapist, dietician, and/or respiratory therapist, depending on local practice is recommended.

Management: Aspiration risks are addressed through careful (e.g., small bites, well-chewed, sipping water between bites of dense foods) or postural feeding, dilation for symptomatic esophageal strictures, GER treatment, and surgical repair of recurrent TEF or laryngeal cleft.

3. Recurrent Infections and Bronchitis:

Complications: Patients with EA-TEF are prone to recurrent lower respiratory infections (pneumonias) and chronic bronchitis due to impaired airway clearance caused by tracheomalacia and recurrent aspiration.

Diagnosis: The diagnosis of bronchitis is based primarily on symptoms. A diagnosis of pneumonia or atelectasis should be confirmed with chest X-ray and/or ultrasound. When recurrent pneumonia is documented, a CT scan is recommended to identify bronchiectasis and exclude alternative diagnoses. Sputum or bronchoalveolar lavage (BAL) can help detect airway colonization by bacteria.

Treatment: start age-appropriate airway clearance techniques to aid in the removal of mucus (in adolescents/ adults: active cycle of breathing techniques) such as positive expiratory pressure devices like PEP masks or oscillatory positive pressure devices^{2,3}. Oral antibiotics – usually for 10-14 days – are recommended when infection is strongly suspected or confirmed, and azithromycin may be used preventatively due to its anti-inflammatory and antibiotic properties.

4. Atelectasis

Pathogenesis: Endobronchial secretions and/or aspiration may lead to atelectasis (areas of lung collapse). This is a particular problem with tracheobronchomalacia, which impairs mucociliary clearance. Atelectasis can also occur after surgery.

Diagnosis: In cases of new or worsening or recurrent respiratory symptoms, atelectasis should be suspected and imaging (lung ultrasound, chest X-ray, thoracic MRI and/or CT scan) should be performed to exclude or confirm the diagnosis.

Treatment: Conservative therapy may include airway clearance techniques, positive airway pressure devices and/or high-flow or CPAP support, use of hypertonic saline nebulisation, mucolytic agents, and eventually systemic steroids, and systemic antibiotics. In persistent atelectasis, bronchoscopy – preferably within 4-6 weeks – may be required to reopen the lung and prevent permanent scarring and bronchiectasis. Follow-up imaging is required to confirm that atelectasis has resolved.

5. Bronchiectasis Monitoring:

Definition: Bronchiectasis, a serious and potentially permanent complication, involves the floppiness and dilation of the airways caused by recurrent and/or persistent bacterial infection and aspiration, leading to chronic bacterial infection.

Diagnosis: When bronchiectasis is suspected, it should be confirmed with a CT scan or MRI of the chest. Depending on the clinical course, annual chest X-rays can be considered to monitor for bronchiectasis, likewise, regular lung function testing.

Management: Early detection, regular treatment with airway clearance techniques, nebulization of hypertonic saline, long-term use of azithromycin, and prompt use of antibiotics during exacerbations can help slow the progression of bronchiectasis.

6. Chronic Respiratory Symptoms (Wheezing, Coughing):

Diagnosis: Symptoms such as wheezing are, after TEF, often caused by airway collapse (TBM) rather than asthma. Misdiagnosis can lead to unnecessary treatments. Nevertheless, a diagnosis of asthma needs to be excluded.

Treatment: Bronchodilators are generally not recommended in case of TBM as they may worsen airway collapse, but they can be trialed for cases with airway hyperreactivity. In infants and preschool children, risk factors (eg. family history of atopic diseases, eczema) and allergy testing may identify children with co-existent asthma. Pulmonary function testing, including bronchodilator response, could additionally be used in individuals 5-6 years and older.

7. Routine Health Maintenance:

Regular Follow-up: Patients should be monitored by a multidisciplinary team (which usually include a pediatric pulmonologist, gastroenterologist, surgeon and/or ear, nose and throat surgeon), with at least annual evaluations.

Vaccinations: In addition to routine vaccinations, annual influenza vaccination is strongly recommended. RSV vaccination should be performed in infants, in accordance with local guidelines.

Lifestyle and Care: Parents and patients should be educated on airway clearance, recognizing early signs of infection, identifying changes in their respiratory status, monitoring for signs of aspiration, and the importance of long-term follow-up care.

References

- 1 Koumbourlis, et al. Care recommendations for the respiratory complications of esophageal atresia-tracheoesophageal fistula. *Pediatr Pulmonol* 2020;55:2713-2729.
- 2 Chang, et al. European Respiratory Society guidelines for the management of children and adolescents with bronchiectasis. *Eur Respir J.* 2021;58:2002990.
- 3 Herrero-Cortina, et al. European Respiratory Society statement on airway clearance techniques in adults with bronchiectasis. *Eur Respir J.* 2023;62:2202053.

This document is endorsed by the Esophageal ATresia global support groups (EAT), the European Reference Network for rare Inherited and Congenital Anomalies (ERNICA), the European Reference Network for rare respiratory diseases (ERN-LUNG), and the International Network of Esophageal Atresia (INoEA).